

Planet Holloway websheet 5.1

AP Physics C - Chapter 5

You may print this out and write on it or work on your own paper.

Show all work.

1. A horizontal force of 120 N is applied to a 22 kg monkey across an 8 m level surface. If the monkey accelerates at 3.5 m/s/s, then what is the work done by the force of friction as it acts to retard the motion of the monkey?
2. Physics Phil pushes a 65 N crate of watermelon up a ramp 16 m in length and inclined at 15 degrees. What is the change of potential energy in the crate as it moves from the bottom of the ramp to the top?
3. A simple pendulum constructed in the front of a physics classroom is 3 m long and has a bowling ball as a bob with a mass of 12 kg. If the support string is initially 25 degrees from vertical and released with an initial speed of 2 m/s, what is the maximum angle it will swing to on the other side?
4. Captain Hooke has a Hooke's law spring hanging from his hook. If the spring is stretched 17 cm and has 84 J stored in it at this point, what is the spring constant for Hooke's Hooke's law spring upon his hook? (I know you're as hooked on physics as I am).
5. Tico pushes a wheelbarrow filled with peat, weighing 280 N, to the top of 40 m long street inclined at 15 degrees. Unfortunately, after leaving the wheelbarrow at the top of the hill, a strong breeze causes the wheelbarrow to slide back down the hill, during which a 57 N friction force acts on it. What is the kinetic energy of the wheelbarrow at the bottom of the hill?
6. A skydiver (55 kg) with a broken foot jumps out of an airplane at a height of 850 m above the ground. If he lands with a speed of 3.5 m/s, what is the energy that friction removed?

Answers:

- | | |
|-----------------|-------------------------|
| 1. - 344 J | 4. 5813.1 N/m |
| 2. 269.2 J | 5. 618.8 J |
| 3. 32.9 degrees | 6. 4.67×10^5 J |